

## TA 10-Year Stereo Composition Measurement

---

**Douglas R. Bergman\*** and **T.A Stroman**, for the **Telescope Array Collaboration**<sup>†</sup>

*Dept. of Physics & Astronomy and High Energy Astrophysics Inst., University of Utah*

*E-mail:* [bergman@physics.utah.edu](mailto:bergman@physics.utah.edu)

The nuclear composition of primary ultra-high energy cosmic rays (UHECRs) can be inferred from the observed distribution of depths of shower maximum ( $X_{\max}$ ) of the induced extensive air showers. The observed  $X_{\max}$  distributions at various primary energies can be compared with the distributions predicted by detailed detector simulations for any assumed primary particle type and high-energy hadronic interaction model. In this paper, we present measurements of  $X_{\max}$  by the Telescope Array (TA) fluorescence detectors with stereoscopic shower reconstruction using 10 years of data. We find that for all hadronic models considered, the data collected is consistent with a chiefly light UHECR composition.

*36th International Cosmic Ray Conference -ICRC2019-  
July 24th - August 1st, 2019  
Madison, WI, U.S.A.*

---

\*Speaker.

<sup>†</sup>for collaboration list see PoS(ICRC2019)1177

## 1. Introduction

The flux of Ultra-high-energy cosmic rays (UHECRs;  $E > 10^{18}$  eV) is very low, requiring large indirect-detection experiments using the Earth’s atmosphere as a calorimeter. The Telescope Array (TA) Experiment covering 700 km<sup>2</sup> of the desert in central Utah, USA, is the largest UHECR detector in the northern hemisphere [1].

One of the primary objectives of TA is the measurement of UHECR nuclear composition. Extensive air showers (EASs) are produced by “primary” UHECRs incident on the atmosphere. An EAS of a given (primary) energy reaches its maximum size at a slant depth  $X_{\max}$  that gets smaller as the nuclear mass of the primary UHECR gets larger. The TA composition measurement uses the observed  $X_{\max}$  distribution in comparison with the distributions predicted by detailed simulations that accurately model the detector, for various high-energy hadronic interaction models. The high-energy interaction models are uncertain due to extrapolations from measurements at much lower energies.

We present the observed  $X_{\max}$  distribution from over 11 years of TA operation compared with simulated distributions created using an identical data analysis. The CORSIKA p0ogram [2] is used to simulate the longitudinal development of hydrogen and iron primaries using several different models (QGSJET-II-04 [3], QGSJET-II-03 [4], QGSJET-01c [5], EPOS LHC [6], and SIBYLL 2.1 [7]). The detector response is then simulated for each shower and simulated data is produced in the same format as the real data from the detector. The  $X_{\max}$  measurements are performed using shower axes determined by stereoscopic measurements, detailed in Section 2. The detector simulation is described in Section 3, followed by the presentation results in Section 4 and a discussion in Section 5.

## 2. Stereo Analysis

TA consists of three sets of fluorescence detector (FD) telescopes, located 20–30 km apart, overlooking an array of surface detectors [8, 9]. Two of the FD stations employ identical, FADC-based telescope electronics, while the third station uses refurbished sample-and-hold equipment from the High Resolution Fly’s Eye Experiment [10]. The FDs operate only on clear, moonless nights and observe the longitudinal development of EASs. When two FD sites observe the same shower, the intersection of shower-detector planes uniquely determines the location and orientation of the shower trajectory to a high degree of accuracy. If three FD sites observe the same shower, an algorithm selects the best pair of sites based on plane-crossing angles.

Using this shower geometry, and the current atmospheric density profile, one can determine the slant depth along shower track which is observed by each FD pixel. We determine  $X_{\max}$  by an inverse Monte-Carlo (IMC) method, in which the parameters of a Gaisser-Hillas shower-shape [12] are varied to find the shower profile that minimizes a  $\chi^2$  comparison between observed and simulated signals.

Energy reconstruction is done by integration of the best-fit Gaisser-Hillas profile, weighted by a self-consistent energy-deposit model, to obtain a calorimetric energy. A correction for the “missing” energy (from the muons and neutrinos in the shower) is calculated from an analysis of QGSJET-II-03 protons.

An independent reconstruction of the shower profile is done for each FD observing an event using the stereo geometry. This can result in up to three separate measurements of the Gaisser-Hillas parameters for each shower. When two or three measurements all pass profile quality cuts, we use the unweighted average values of  $X_{\max}$  and  $\log_{10}(E/\text{eV})$  in the analysis. If only one measurement survives the cuts, it is admitted to the final data set only if it passes an additional quality cut based on a pattern-recognition analysis originally developed for “hybrid” reconstruction and described in [13].

### 3. Simulation Procedure

We use a shower library consisting of Gaisser-Hillas fits to CORISKA-generated shower longitudinal data as the input to our detection simulation. The shower library has a number of bins in energy and shower inclination. Shower trajectories are selected from an isotropic distribution with zenith angles  $\theta \leq 80^\circ$ . Shower energies are chosen according to the HiRes spectrum [10] for energies above  $10^{17.7}$  eV. The response of the detector is then simulated, including fluorescence and Cherenkov light production, atmospheric transmission, optical acceptance, and detector electronics simulation including night-sky background noise. The detector simulation is performed for all nights when at least two FDs were operating from 4 Nov 2007 to 28 Nov 2017.

The simulation outputs artificial raw data, which can be processed and analyzed with the analysis chain which is applied to natural night-sky data. This chain includes identification of shower-detector planes, inter-FD coincidence detection, stereo geometry calculation, and profile reconstruction [11].


### 4. Results

We show our  $X_{\max}$  distributions in energy bins for  $E \geq 10^{18.4}$  eV and the QGSJET-II-04 predictions for hydrogen and iron in Figure 1. In Figure 2 we show the mean of the observed and simulated distributions in several energy bins, and linear fits to these values from all physics models to illustrate the relationships among the various predictions.


### 5. Discussion and Conclusions

Although iron is an attractive candidate for acceleration to ultra-high energies because of its cosmic abundance, nuclear stability, and large electric charge, the results shown in Section 4 demonstrate that the  $X_{\max}$  distribution observed by stereo analysis of TA data does not support a significant amount of iron in the composition at any energy above  $10^{18.4}$  eV, regardless of the high energy interaction model. A pure-proton composition is attractive when combined with QGSJET-01c, but the agreement is less clear when post-LHC models are considered. This applies marginally to QGSJET-II-04, and more strongly to EPOS LHC, and also to the pre-LHC model SIBYLL 2.1; LHC corrections to SIBYLL are expected to further widen the difference from TA data [14].

On average, our reconstruction of  $X_{\max}$  and energy are respectively accurate to better than 25 g/cm<sup>2</sup> and 7%. The systematic uncertainty on these TA  $X_{\max}$  measurements is approximately 15



**Figure 1:** The distribution of reconstructed  $X_{\max}$ , binned by reconstructed primary energy. The proton and iron predictions are based on the QGSJET-II-04 model for high-energy hadronic interactions. The data and proton histograms agree in both mean and overall shape, while disagreeing strongly with iron.



**Figure 2:** A scatter plot of  $X_{\max}$  vs. energy for 10 years of data, overlaid with profile histograms showing the energy evolution of the mean  $X_{\max}$  of the data and each Monte Carlo prediction (proton and iron, according to QGSJET-II-04). The systematic uncertainty on reconstructed  $X_{\max}$  is  $15 \text{ g/cm}^2$ .

$\text{g}/\text{cm}^2$ . Much of this uncertainty originates in the atmospheric models used, both for the density profile and the aerosol distribution.

Further work on this analysis, will help to clarify the extent to which the UHECR composition can be explained only one light element. Contributions from other nuclear constituents may help to bring various moments of the observed and simulated  $X_{\text{max}}$  distributions into agreement, but we are also exploring the use of statistically robust comparisons that consider the entire distribution. In particular, the Cramér-von Mises test statistic [15], combined with the value of whatever artificially imposed offset minimizes it, shows great promise for quantifying the role played by intermediate-mass elements in the UHECR flux.

## 6. Acknowledgments

The Telescope Array experiment is supported by the Japan Society for the Promotion of Science (JSPS) through Grants-in-Aid for Priority Area 431, for Specially Promoted Research JP21000002, for Scientific Research (S) JP19104006, for Specially Promoted Research JP15H05693, for Scientific Research (S) JP15H05741, for Science Research (A) JP18H03705 and for Young Scientists (A) JPH26707011; by the joint research program of the Institute for Cosmic Ray Research (ICRR), The University of Tokyo; by the U.S. National Science Foundation awards PHY-0601915, PHY-1404495, PHY-1404502, and PHY-1607727; by the National Research Foundation of Korea (2016R1A2B4014967, 2016R1A5A1013277, 2017K1A4A3015188, 2017R1A2A1A05071429) ; by the Russian Academy of Sciences, RFBR grant 16-02-00962a (INR), IISN project No. 4.4502.13, and Belgian Science Policy under IUAP VII/37 (ULB). The foundations of Dr. Ezekiel R. and Edna Wattis Dumke, Willard L. Eccles, and George S. and Dolores Doré Eccles all helped with generous donations. The State of Utah supported the project through its Economic Development Board, and the University of Utah through the Office of the Vice President for Research. The experimental site became available through the cooperation of the Utah School and Institutional Trust Lands Administration (SITLA), U.S. Bureau of Land Management (BLM), and the U.S. Air Force. We appreciate the assistance of the State of Utah and Fillmore offices of the BLM in crafting the Plan of Development for the site. Patrick Shea assisted the collaboration with valuable advice on a variety of topics. The people and the officials of Millard County, Utah have been a source of steadfast and warm support for our work which we greatly appreciate. We are indebted to the Millard County Road Department for their efforts to maintain and clear the roads which get us to our sites. We gratefully acknowledge the contribution from the technical staffs of our home institutions. An allocation of computer time from the Center for High Performance Computing at the University of Utah is gratefully acknowledged.

## References

- [1] J. Matthews *et al.*, *Overview of the Telescope Array Experiment*, Proceedings of the 31st ICRC (Łódź), p. icrc1386, 2009.
- [2] D. Heck *et al.*, *CORSIKA: A Monte Carlo Code to Simulate Extensive Air Showers*, FZKA 6019 (1998).

- [3] S. Ostapchenko, *QGSJET-II: physics, recent improvements, and results for air showers*, in European Physical Journal Web of Conferences, vol. 52 of European Physical Journal Web of Conferences, p. 2001, June, 2013.
- [4] S. Ostapchenko, *QGSJET-II: towards reliable description of very high energy hadronic interactions*, Nuclear Physics B Proceedings Supplements **151** (2006) 143. [hep-ph/0412332]
- [5] N.N. Kalmykov, S.S. Ostapchenko, and A.I. Pavlov, *Quark-Gluon-String Model and EAS Simulation Problems at Ultra-High Energies*, Nuclear Physics B Proceedings Supplements **52** (1997) 17.
- [6] T. Pierog *et al.*, *EPOS LHC : test of collective hadronization with LHC data*, arXiv:1306.0121.
- [7] E.-J. Ahn *et al.*, *Cosmic ray interaction event generator SIBYLL 2.1*, Phys. Rev. D **80** (2009) 094003. [arXiv:0906.4113]
- [8] H. Tokuno *et al.*, *New air fluorescence detectors employed in the Telescope Array experiment*, Nuclear Instruments and Methods in Physics Research A **676** (2012) 54. [arXiv:1201.0002]
- [9] T. Abu-Zayyad *et al.*, *The energy spectrum of Telescope Array's Middle Drum detector and the direct comparison to the High Resolution Fly's Eye experiment*, Astropart. Phys. **39** (2012) 109. [arXiv:1202.5141]
- [10] R.U. Abbasi *et al.*, *First Observation of the Greisen-Zatsepin-Kuzmin Suppression*, Phys. Rev. Lett. **100**(2008) 101101. [astro-ph/0703099]
- [11] T. Abu-Zayyad *et al.*, *The energy spectrum of ultra-high-energy cosmic rays measured by the Telescope Array FADC fluorescence detectors in monocular mode*, Astropart. Phys. **48** (2013) 16. [arXiv:1305.6079]
- [12] T.K. Gaisser and A.M. Hillas, *Reliability of the method of constant intensity cuts for reconstructing the average development of vertical showers*, in International Cosmic Ray Conference, vol. 8 of 15th ICRC (Plovdiv), pp. 353–357, 1977.
- [13] R.U. Abbasi *et al.*, *Study of Ultra-High Energy Cosmic Ray composition using Telescope Array's Middle Drum detector and surface array in hybrid mode*, Astropart. Phys. **64** (2015) 49. [arXiv:1408.1726]
- [14] E. Ahn *et al.*, *LHC Update of the Hadronic Interaction Model SIBYLL 2.1*, in International Cosmic Ray Conference, 33rd ICRC (Rio de Janeiro) abstract 0803, 2013.
- [15] T.W. Anderson, *On the Distribution of the Two-Sample Cramer-von Mises Criterion*, Ann. Math. Statist. **33** (1962) 1148.